# COVID-19 Vaccine Timing and Utilization

## Initial Vaccine Doses

**Pfizer-BioNTech Vaccine (30mcg x 2 doses 21 days apart: ≥12yo; 10mcg x 2 doses 21 days apart: 5-11yo)**

→ **≥ 28 days** → **Pfizer-BioNTech** Moderately to severely immunocompromised individuals (≥5yo) → **Booster ≥ 5 Months** → **Pfizer-BioNTech Dose: 30mcg**
- ≥ 12yo

→ **Booster ≥ 5 Months** → **Pfizer-BioNTech Dose: 30mcg**
- ≥ 12yo

**Moderna Vaccine (100mcg x 2 doses 28 days apart)**

→ **≥ 28 days** → **Moderna** Moderately to severely immunocompromised individuals (≥ 18yo) → **Booster ≥ 5 Months** → **Moderna Dose: 50mcg**
- ≥ 18yo

→ **Booster ≥ 5 Months** → **Moderna Dose: 50mcg**
- ≥ 18yo

**Janssen (J&J) Vaccine (0.5 mL single dose)**

→ **Booster ≥ 2 Months** → **Pfizer or Moderna booster preferred in most situations. If using Janssen:**

**Janssen (J&J) Dose: 0.5 mL**
- ≥ 18yo

## Additional Dose

## Booster Dose

January 2022 Interim Clinical Considerations for Use of COVID-19 Vaccines | CDC

Kentucky Public Health
Prevent. Promote. Protect.